

SATELLITE INTENSITY PREDICTIONS OF  
NORTH PACIFIC CYCLOGENESIS

Waldo J. Younker Jr.  
NATIONAL ENVIRONMENTAL SATELLITE SERVICE  
ANCHORAGE ALASKA 99513

1. Introduction

The NOAA-5 and GOES infrared (IR) satellite imagery received at the Anchorage SFSS, called to my attention a particular cloud pattern associated with some of the most rapid developing cyclogenesis cases over the Pacific. I will refer to this particular cloud pattern as an S-C cloud system. It has been noted that these S-C systems almost invariably result in deepening surface low pressure centers, as long as the center stays over water and does not get captured by an upper low center. This fact seems to be almost independent of whether or not the center of the S-C system is behind or in advance of the 500-mb trough.

I thought of using this recurring satellite observed cloud pattern in conjunction with deepening prediction graphs once used in the Anchorage forecast office. The prediction graphs, published in the July 1959 AMS Bulletin by Snopkowski and Welch(1), were frequently used to help make hand drawn subjective surface prognostic charts. The prediction graphs had fallen into relative disuse by the 1970's, well before a steady flow of IR satellite imagery started here in 1976.

The objective surface low center deepening method used at Anchorage in the 1960's followed the conclusions in the Bulletin article, which recommended using for the Western Pacific, only the modal values from the earlier Project-AROWA(2) Atlantic graph (fig. 1), along with a modified Eastern Pacific graph (fig. 2).

The criteria for using these two graphs were:

1. Use between 30N to 60N latitude only.
2. The 180th meridian separates East from West Pacific.
3. For surface lows where centers are located under the 500-mb contours between the trough and down stream ridge, forecast decreasing central pressure.
4. For lows located directly under the 500-mb trough - i.e., within one degree longitude on either side, or located under the forward half (with respect to movement) of the closed 500-mb contours, forecast no change in central pressure.
5. For lows located elsewhere, forecast increasing surface central pressure.
6. Do not use these graphs for tropical storms which are in transition from typhoons to extratropical cyclones.




Figure 1. North Atlantic deepening prediction graph. Use only modal values for Western Pacific. (Snopkowski and Welch)




Figure 2. Deepening Prediction graph for Eastern Pacific, mb per 24 hr (Snopkowski and Welch)

## Method of computation:

Measure the 500-mb height difference between points 7.5 degrees of latitude either side of the surface low position normal to the 500-mb contours. When closed contours lie in this range, measure only to the last open contour (fig. 3).




Figure 3. Method for determining 500mb height difference for computation. (Snopkowski and Welch)

Measure the 500-mb temperature difference from the surface low position within the sector extending 15 degrees of latitude northwest of the low (fig. 4).

With the above two parameters, use either the Western Pacific deepening-prediction graph (fig. 1) or the Eastern Pacific graph (fig. 2), or a combination of both, in order to obtain the 24 hour deepening prediction.




Figure 4. Method for determining 500mb temperature difference for deepening computation. (Snopkowski and Welch)

To use these deepening-prediction graphs, the forecaster had to be aware of the existence and position of a surface low pressure center along with its central pressure. In addition, the Anchorage forecasters found that they did not feel secure enough to rely solely on this technique. The statistical results in the Bulletin article indicated deepening forecast cases for the Western Pacific were overforecast by 10 percent in the winter and 33 percent in the summer, and overforecast for the Eastern Pacific 7 percent in the winter and 9 percent in the summer. The added assurance needed to avoid this overforecasting by Anchorage forecasters was the surface evidence of deepening in successive analyses. Consequently, computes were not frequently used on surface lows in their initial stages of development.

Since the often observed S-C cloud pattern seemed to identify the position of impending cyclogenesis up to 12 hours before a low was analyzed at the surface, and thus well before significant deepening had occurred, it was hoped that a combination of these deepening-prediction graphs with S-C cloud patterns would give objective forecasts up to 24 hours earlier than previously obtainable. The purpose of this paper is to show the encouraging results of this marriage.

## 2. Identification of S-C cloud systems and their associated cloud circulation center positions

The diagrams (fig. 5) are what I refer to as S-C cloud structures. The letters S and C refer to the cloud shape on the cold side of the baroclinic zone or wind maximum. Instead of looking at upward vertical motion induced cloudiness, I focus more attention in these cases on the cloud structure that appears to be a direct result of sinking motion aloft. It seems that this sinking motion over or to the rear of a deep cloud circulation center is a more timely clue for cyclogenesis over the North Pacific.

In both the S and C cases sketched in Figure 5, the low clouds to the right of the dashed lines tend to be near the same temperature (same shade of gray on the Z curve display) as that of the underlying water. Consequently, it is often difficult, in Z curve imagery, to distinguish whether this area is clear or covered by a low overcast. The southern end (point L) of the S and C cloud structure is also frequently indistinguishable, since these clouds are low. There must be a gradual increase in brightness from the southern or lower portion of the S or C cloud structure to the northern or upper portion (L to U) without a leaf or layered appearance. Additionally, clouds in the L and U band of Figure 5a become colder towards the western edge. These features can be seen in the IR pictures of Figures 6 and 7.




Figure 5. Schematic showing "S" and "C" cloud patterns.




Figure 6. GOES-3 IR imagery 1145 GMT 1 October 1978.



Figure 7. GOES-3 IR imagery 0745 GMT 12 October 1978.

It is known that the S stage is an early stage in the development of circulation. The C stage is a later stage which occurs when the rotational component of the wind around the low level circulation center becomes faster than the speed of translation of the circulation center with respect to the earth's surface. At this time when return flow begins around the rear of the cloud system, the southern portion (L) of the S cloud band turns to become the cyclonically curved southern portion of the C cloud. Since these low level cloud tops are often near or below the 850-mb level, they are a good indication that a surface wave with open isobars has changed into a closed low once the C cloud stage is reached.

The "X's" (fig. 5) represent the most frequent observed position of the lower cloud circulation centers. In case b, this lower cloud center can most often be located by imagining the continuation of the curvature of the inside of the dark cloud band (L to U) around in an oval shape under the PVA comma tail (indicated by dots). The center of this oval is the position (X) of the lower cloud circulation center. Notice that this point is well to the rear of the upper cloud vortex "V" which is the position

of the upper cloud circulation center. Of all strong cyclogenesis cases over the North Pacific, this large slope by as much as 200 miles between upper and lower circulation centers seems to be restricted to this C cloud pattern. It is important to use this lower cloud circulation position in these C type cloud cases to obtain the best estimate of the surface low position.

Experience has shown that a definite relationship exists between positions of cloud circulation centers and surface lows over the Pacific from 25N to 65N during the cooler months of October through May. Surface low pressure centers will be about one degree of latitude to the left of the direction of translation of significant cloud circulation centers with a 30 kt speed of movement, and within 2 degrees latitude at 60 knots. This relationship is the same for both the more common upper cloud circulation centers that reflect down to the surface without exposing a middle or lower cloud center, and the lower cloud circulation centers of the S-C type systems. This rule does not work well over the North Pacific from June through September apparently due to the lack of enough vertical extent in the circulation patterns.



The accuracy in determining the surface low pressure center position from S-C cloud patterns using this displacement rule is dependent on skill in finding the lower cloud circulation center position. Even though the easier to see upper cloud circulation center in IR pictures will give the general location, the better related lower cloud circulation center is often hidden in more uniform dark grey low clouds. When a visual picture is available, the location of the lower cloud circulation center is often simplified since high clouds are usually not masking this center. A good example of this problem appears in the left hand portion of the UC-2 sector of Figure 8. Test yourself by looking at this IR picture

and guessing what part, if any, is free of clouds in the vicinity of this storm, and also try to estimate the position of the lower cloud circulation center. Then refer to the visual picture (fig. 9) of this same storm to find the answers. This extra-tropical low is experiencing frontogenesis just prior to intense deepening. Note that the clouds associated with the PVA comma (P) in Figure 8 slope down on the inside edge, in a funnel shape, to the lower cloud circulation center apparent in Figure 9. This evidence of sinking air results in the same shading (change in brightness on the inside edge of the high clouds) as that normally seen in the S and C clouds bands previously discussed.




Figure 8. GOES-3 IR imagery 2245 GMT 5 October 1978.




Figure 9. GOES-3 Visible imagery 2315 GMT 5 October 1978.

### 3. Technique of making 24-36 hour surface low central pressure predictions

After tracking an S-C type cloud circulation center to obtain the speed and direction of translation for at least 8 hours, the previously mentioned displacement rule was used to estimate corresponding surface positions of the impending or already formed and intensifying low center. After obtaining the surface pressure from an NMC or Anchorage WSFO analysis at a 00 GMT or 12 GMT position along with the corresponding NMC 500-mb temperature and height parameters (figs. 3, 4), one or both the deepening-prediction graphs (figs. 1, 2) were used to arrive at a final 24 or 36 hour objective central pressure prediction. In cyclogenesis cases in which a surface low was expected to remain over water and not become captured by an upper low or move through a long wave upper ridge position, the forecasts were extended to 36 hours using the 24 hour deepening rates indicated by the graphs.

### 4. Results of first 40 predictions using only S-C cloud centers

Of the 40 deepening predictions made through November 10, 1978, the last 36 were issued in Satellite Interpretation Messages (SIM's) well in advance of their verifying time.

<u>computes</u>	<u>results</u>
32	within 1 mb
5	within 2 mb
1	underforecast 6 mb, but 8 mb better than the LFM forecast
1	underforecast 10 mb, but 10 mb better than the LFM forecast
1	non-deepening case

Several of them were extended forecasts for 36 hours; however the lead time in the SIM's were only 15 to 27 hours primarily due to the 7 hour delay in reception of the NMC 500-mb analysis. Thirty-nine of these 40 predictions were based on the Eastern Pacific graph. Of these 40 cases, 39 deepened and verified better than the latest machine progs that were available at the time the computes were made. The only non-deepening case was one of earliest attempts in this technique. This error was due to a premature classification of the cloud system in the distorted area at the picture edge. When this cloud system moved into better view just 2 hours after the SIM had been transmitted, it was then obvious that this was not the S-C cloud pattern it was previously thought to be. If this one mistake were thrown out, the frequency of deepening indicated by this technique would have been 100 percent instead of just 98 percent.

It was found after verification that the forecast depth of the low centers was within 2 mb of the actual central pressure in 93 percent of the 40 predictions. If the non-deepening case were disregarded, 95 percent of the computes were within 2 mb of the actual central pressure. The results through November 10, 1978 were:

The following example is one of these extended forecast cases that also needed to be interpolated between the 24-hour and extended 36-hour deepening predictions from the graphs, because the S-C cloud center was expected to deepen beyond 24 hours but only remain offshore 33 hours from its 00 GMT position. This information was included in the following portion of the October 14, 1978 SIM.

ANC W TBXX6 PANC 140900 SFSS ANC SIM.  
IMAGERY THRU 0745Z

WIND MAXIMUM WELL DEFINED NORTHEAST PACIFIC BY SHARP EDGE TO PVA SOUTHEAST OF WIND MAXIMUM NOW ALONG ARC 50/152-59/145-61/135 EASTSOUTHEAST. NO BUILDING NORTHWARD OVER SOUTHEAST ALASKA YET APPARENT THE PAST 6 HOURS ALTHOUGH MOISTURE IS INCREASING, WELL DEFINED SHORT WAVE TROUGH UPSTREAM SOUTHEAST ACROSS THE BAROCLINIC ZONE 43/152-39/142 MOVING NORTHEAST 40-45 KNOTS. PROBABLE SUR-

FACE LOW ABOUT 90 MILES NORTHWEST OF APPARENT LOW CLOUD CIRCULATION CENTER 42/148. DEEPENING GRAPH GIVES 3-4MB DEEPENING IN 24 HOURS FROM 00Z...THUS EXPECT ABOUT 1004MB CENTER AT 00Z TOMORROW ENTERING THE SOUTHEAST PORTION OF THE GULF OF ALASKA. EXTENDING THE TRACK AT THE PRESENT RATE OF MOVEMENT GIVES 51/140 POSITION FOR THE CIRCULATION CENTER...THUS A SURFACE LOW JUST NORTHWEST NEAR 52/141 IS FIRST GUESS FOR 00Z. THIS IS VERY CLOSE TO THE POSITION OF AN INVERTED SURFACE TROUGH ON LFM AT THAT TIME...BUT 12MB DEEPER. FASTER DEEPENING EXPECTED AS FORECAST BY LFM THEREAFTER UNTIL THE LOW HITS THE COAST LINE...THUS CENTRAL PRESSURE OF 8MB DEEPER THAN LFM AND PE LIKELY WHEN THE CENTER IS EXPECTED TO HIT SOUTHEAST ALASKA COAST ROUGHLY 09Z...AND POSSIBLY 12MB DEEPER. THIS COMES OUT TO A 996MB TO 1000MB CENTER TO HIT COAST WHEN TRYING TO EXTEND THE DEEPENING GRAPH PREDICTION OUT BEYOND THE 24 PERIOD IT WAS DESIGNED FOR.



